

Classifying landform at broad spatial scales: the distribution and conservation of wetlands in New South Wales, Australia

R. T. Kingsford^A, K. Brandis^A, R. F. Thomas^A, P. Crighton^B, E. Knowles^A and E. Gale^C

^ANew South Wales Department of Environment and Conservation, PO Box 1967, Hurstville, NSW 2220, Australia.

^BAustralia Water and Power Pty Ltd, Level 6, 185 Macquarie Street, Sydney, NSW 2000, Australia.

^CCentre for Water Research, University of Western Australia, 35 Stirling Highway, Crawley, WA 6009, Australia.

^DCorresponding author. Email: richard.kingsford@npws.nsw.gov.au

Abstract. Relatively few large-scale inventories of the world's wetlands exist because of the difficulties of spatial scale, associated cost and multiple objectives, often temporally confounded, that drive classification. The extent of wetlands across a large part of Australia (New South Wales, 80.6 million ha) was determined using satellite image analyses. These data allowed analyses of the distribution of wetlands, their conservation status and potential threats at different spatial scales; that is, State, coastal and inland, and catchment. Approximately 5.6% of New South Wales is wetland (4.5 million ha), mostly (96%) in inland river catchments. Broad classification allowed identification of the extent of wetland types: (i) floodplains (89%); (ii) freshwater lakes (6.6%); (iii) saline lakes (< 1%); (iv) estuarine wetlands (2.5%); and (v) coastal lagoons and lakes (1.5%). Conservation reserves protect only 3% of wetland area. The analyses identified the north-west as the key area for wetland conservation as most other catchments have lower wetland extent and more potential threatening processes. The first stage of a large-scale inventory is to determine the extent and location of wetlands, with immediate benefits for strategic conservation and management. Other objectives (e.g. classification, biotic composition, hydrology and threats) seldom have sufficient data available for large-scale inventories but can be completed later with resources.

Extra keywords: catchments, classification, inventory, mapping, reserves, river regulation.

Introduction

Wetlands are among the most threatened ecosystems in the world (Lemly *et al.* 2000) and Australia is no exception (Finlayson and Rea 1999). The major factor implicated in their destruction is water resource development and the draining of wetlands (Kingsford 2000a), although urbanisation in coastal areas is significant (Adam 1995). Wetlands also support high levels of biodiversity and have important cultural and economic values. There is increasing international commitment to wetland conservation, with many countries signing the Convention on Wetlands (Ramsar Convention) but objective information is needed for effective conservation.

It is difficult to manage any natural resource without knowing its distribution, and wetlands are no exception (Finlayson *et al.* 1999). Large-scale inventories are necessary for conservation management and effective planning (Pressey and Adam 1995). Signatories to the Ramsar Convention resolved to develop low-cost and user-friendly methodology for inventories (The Ramsar Convention

Bureau 1999). Of existing inventories of wetlands, relatively few adequately define objectives, determine the distribution and size of wetlands, provide explicit methodology, differentiate wetland area according to classification or are publicly accessible (Finlayson *et al.* 1999). The notable exception is the large-scale inventory of wetlands in North America (Wilen and Bates 1995). Producing inventories of wetlands remains elusive, constraining effective management of these biodiverse ecosystems.

Part of the problem is attributable to multiple objectives for inventories. These may include identifying location and extent, geomorphology, soil type, water regimen, water chemistry, biota, anthropogenic threats and management regimen (see core essential data elements and additional categories in Finlayson *et al.* 1999). These represent a mix of spatial and temporal variables that are difficult to collect over large spatial scales. Some are seldom available (e.g. water regimen, chemistry or biota), except for small areas. A large-scale inventory should initially determine the extent and distribution of wetlands and this can be done with remote

sensing at large spatial scales, although trade-offs between resolution and budget exist (Wilén and Bates 1995). With the resulting spatial layer of wetlands, a second objective can be achieved; that is, analysis of the spatial distribution of wetlands at different scales (e.g. catchments), in relation to conservation effort or threats.

Classification of wetlands is not necessary for such analyses but it adds value in understanding representativeness. It remains a vexed question for inventories (Pressey and Adam 1995; Wilén and Bates 1995; Semeniuk and Semeniuk 1995, 1997) mainly because all classifications are simple representations of considerable spatial and temporal complexity. The hierarchical classification from North America is used globally (Cowardin *et al.* 1979; Wilén and Bates 1995) and adapted for parts of Australia (Pressey and Adam 1995) but it has problems with definition and lacks exclusivity in categories (Semeniuk and Semeniuk 1995, 1997). A suggested replacement, the hydro-geomorphic classification (Semeniuk and Semeniuk 1995, 1997), simplifies hydrological variability according to permanent, seasonal and intermittent, which are categories seldom known for wetlands across large areas. Ideally, classifications of wetlands should be quantitative and based on the objectives of the classifications that vary (Pressey and Adam 1995; Rempel *et al.* 1997).

Our primary objective was to do a large-scale inventory to identify the extent of wetlands across a large part of Australia (the State of New South Wales (NSW)) because Australia devolves responsibility for natural resource management, including water, to the provincial level of States. Second, we analysed the distribution of wetlands according to elevation, climate and different spatial scales; that is, statewide, coastal and inland regions, and catchments. The spatial framework was extended to an analysis of conservation effort (reserves) and threats to wetlands to provide a foundation for strategic policies for wetland conservation, including river management planning and conservation of aquatic fauna and flora.

Materials and methods

Wetland distribution and extent

For the 17 inland catchments, wetlands were mapped from Landsat Multi-Spectral Scanner (MSS) (80 m pixel) data (40 scenes), whereas the 22 coastal catchments wetlands were mapped from Landsat Thematic Mapper (TM) data (30 m pixel) (11 scenes) (Fig. 1). We chose the wettest times within a 10-year period (1984–1993) for inland areas. These were identified using monthly National Oceanographic and Atmospheric Administration (NOAA) satellite imagery, wetland data from aerial surveys across 10% of the State (Kingsford *et al.* 1999) and annual rainfall within each river catchment. Additional images were chosen for areas where high catchment rainfall did not coincide with high local rainfall. Cloud-free images and a period of 16 days for the imagery determined the actual dates of the images used. For coastal catchments where wetlands were less transient, cloud cover was the primary determinant of image suitability and we used cloud-free imagery within a 6-month period (September 1994 to January 1995). Each Landsat image was rectified and georeferenced to 1:250 000 and

1:250 000 topographic maps for the inland and the coast, respectively, using the Universal Transverse Mercator projection (Australian Map Grid) with 14–30 ground control points, and evenly distributed across each image.

For inland catchments, wetland boundaries were delineated from non-wetland areas using an unsupervised classification (Richards 1993) that was based on the variable spectral signatures for water (open water and inundated vegetation) and for adjacent areas of dependent aquatic vegetation. For coastal catchments, we used a band 5 density slice to delineate open water body boundaries (Frazier and Page 2000) and a supervised classification (ERDAS 2001) to delineate adjacent coastal wetland vegetation. Spectral similarities among wetland and non-wetland areas (Johnston and Barson 1993; Frazier and Page 2000) demanded an iterative process of systematic visual inspection and reclassification for each satellite image to initially validate wetland boundaries (Congalton 2001). This involved draping the wetland boundaries over satellite imagery, followed by comparison with the best available ancillary data. For the inland, this included aerial survey data (Kingsford *et al.* 1999), 1:250 000 topographic map sheets and digital coverage of water bodies (AUSLIG 1994), and for the coastal region, this involved 1:250 000 topographic maps and aerial photography. These data helped discriminate between wetland areas with similar spectral signatures to non-wetland areas (e.g. cultivated crops, flooded urban areas and hillslope shadows).

We classified wetlands *a posteriori* using a combination of spectral information and ancillary data. First, we separated water bodies into the three broad groups, which are recognised by the Ramsar Bureau: coastal, inland and man-made (reservoirs) (Semeniuk and Semeniuk 1997). We then used the broadly accepted major geomorphic groups: estuarine, riverine, lacustrine and palustrine (Wilén and Bates 1995) to classify wetlands but we could not objectively separate palustrine and riverine (Table 1). We used spectral and ancillary data to separate further the lacustrine group into freshwater lakes, saline lakes, and coastal lagoons and lakes (Table 1). A more detailed classification of wetlands was not possible because of the variable spectral signature, particularly of floodplains, and the poor availability of consistent ancillary data across the entire study area.

We derived error matrices of the mapped wetland boundaries and aerial photography reference data (Congalton 1991) using a stratified random sampling for two inland catchments (see Kingsford *et al.* 2001) and stratified random sampling for coastal catchments. This produced two quantitative measures of classification accuracy as a whole, overall classification accuracy (total per cent of correctly mapped polygons; Congalton 1991) and the Kappa statistic (actual agreement between mapped and reference data minus chance agreement; Congalton and Green 1999). Kappa also accounts for errors within individual classes of the error matrix (Rosenfield and Fitzpatrick-Lins 1986). We found an overall accuracy of 86% and a Kappa statistic of 0.76 for inland catchments of the Paroo and Warrego River (Kingsford *et al.* 2001). For the coastal catchments, overall accuracy for the Richmond, the north coast (Tweed to Macquarie–Tuggerah Lakes) and the south coast (Hawkesbury–Nepean to Snowy) were 80% (Kappa = 0.66), 85% (Kappa = 0.80) and 90% (Kappa = 0.87), respectively. Such overall classification accuracies are considered good (> 80%) (Congalton 1991), and the Kappa statistics represented strong agreement (> 0.80) (north coast and south coast) and moderate agreement (> 0.40) (Paroo and Warrego, and Richmond) between mapped data and reference data (Congalton 2001).

We determined the area and proportion of land area of wetland groups and reservoirs within each catchment across inland and coastal catchments. For river catchments within the Murray–Darling Basin across State borders, we analysed wetland distribution for the whole catchment and separately for each State. We compared the distribution (coastal and inland) and extent of wetlands in relation to elevation (above sea level) and mean annual rainfall between 1980 and 1999.




Fig. 1. Locations of 39 catchments in New South Wales (NSW) and the Murray–Darling Basin (grey inset in map of Australia with State borders), including 17 inland catchments of rivers flowing west of the Great Dividing Range (1–17) and 22 coastal catchments of rivers flowing east of the Great Dividing Range (22–39) (grey shading). Numbers for catchments match names given in Tables 3–6.

Wetland conservation

We used GIS analyses to determine the area of wetlands and reservoirs protected under legislative or policy instruments that confer status or protection on wetlands that were applied to the land area of the wetland. These represent the most used and applicable measures for assessing wetland conservation. They included four conservation measures: (i) National Parks and Wildlife Service (NPWS) reserves; (ii) State Environmental Planning Policy No. 14 (SEPP 14) for coastal wetlands (Farrier *et al.* 1999); (iii) the Directory of Important Wetlands in Australia (DIWA) (Environment Australia 2001); and (iv) wetlands listed under the Ramsar Convention. We also included State forest reserves because, although set aside primarily for timber production, some have become ‘quasi’ conservation areas for wetlands (e.g. Barmah–Millewa Forest). The wetland coverage for each catchment was clipped in Arc/Info (ESRI 2000) using digital boundaries for the relevant conservation measures. We also assessed potential threats to wetlands by overlaying the proportion of land in each catchment covered by intensive agriculture (National Land and Water Resources Audit 2001a), number of weirs and storage capacity of dams (Kingsford 1995; National Land and Water Resources Audit 2001b).

Results

Wetlands of New South Wales

There are almost 4.5 million ha of wetlands in NSW or 5.6% of the land area of the State (Table 2, Fig. 2). Most wetland area (96%) is in the inland, in the catchments of the Murray–Darling Basin and far west of the State (Fig. 2), with nearly half of all inland wetland area (46%) in the Far north-west, Paroo, Warrego and Condamine–Culgoa catchments (Table 3). Of the wetland types across the State, floodplain wetlands are the most ubiquitous, forming 89% of all wetland area and covering 5% of the land surface (Table 2). Freshwater lakes and estuarine areas make up most of the remaining wetland area in the State, with comparatively small areas of saline lakes and coastal lagoons and lakes (Table 2). There are more reservoirs than freshwater lakes, coastal lagoons and lakes, and salt lakes.

Most wetland area (> 95%) is below 250 m above sea level, and 93% is in parts of the State that receive less than 500 mm of annual rainfall (Fig. 3).

Inland wetlands

Most inland wetlands are floodplains that are distributed below 250 m and where there is less than 500 mm of annual rainfall (Table 2; Fig. 3). The highest proportion of wetlands in NSW is in two catchments, Warrego and Condamine–Culgoa, whereby 27% and 28% of their respective areas in the NSW part of the catchment is covered by wetlands (Table 3). The Condamine–Culgoa has the highest wetland area of any catchment in inland NSW, covering approximately 740 000 ha (Table 3). The Paroo and Murray–Riverina catchments have the next highest proportion of wetland with 16% and 17% of land area covered, respectively. Most of the wetland area and higher proportions in the Paroo, Warrego and Condamine–Culgoa are on the NSW side of the border, and there are similar proportions for the Lower Murray River catchments and the Border and Moonie Rivers on both sides of the border (Table 3). Nine of the 17 inland catchments in NSW have more than 200 000 ha of wetland and the remainder of the catchments has less than 100 000 ha of wetland (Table 3). Most catchments with lower areas of wetland also have a low proportion of wetlands, with the exception of Lake George catchment, which ranks fifth in terms of wetland proportion. Twelve of the 17 inland catchments each have 6% or less of wetland area in their catchment (Table 3). These include most of the major regulated rivers such as Darling, Border and Moonie, Gwydir, Namoi, Macquarie–Bogan, Lachlan, Murrumbidgee, Upper Murray and Lower Murray (Table 3). The Murray River and its tributaries flow through four of the catchments (Table 3). The Murray–Riverina catchment includes most of the floodplain of this river and when all catchments of the Murray River are combined, the proportion (7%) is similar to that of other regulated river catchments (Table 3).

There is more than 70% of wetland area identified as floodplain in all but two of the catchments and, in 12 catchments (Paroo, Warrego, Condamine–Culgoa, Border and Moonie, Gwydir, Castlereagh, Macquarie–Bogan, Lachlan, Murrumbidgee, Upper Murray, Murray–Riverina and Benanee), there is more than 90% floodplain wetland (Table 3). The Paroo and Condamine–Culgoa catchments have more than half a million hectares of floodplain wetland in NSW, and another seven each have more than 200 000 ha of floodplain (Warrego, Darling, Macquarie–Bogan, Lachlan, Murrumbidgee, Murray–Riverina and Far north-west).

The Darling catchment has the greatest area of freshwater lakes (130 730 ha) of any inland catchment, primarily the Menindee Lakes, and the Far north-west has an order of magnitude of more freshwater lakes (Table 3). Freshwater lake area exceeds 10 000 ha in each of seven other

catchments: Paroo, Condamine–Culgoa, Lachlan, Murrumbidgee, Lake George, Lower Murray and Far north-west. The Castlereagh, Murray–Riverina, Upper Murray, Macquarie, Gwydir, Border and Warrego have little (< 1%) or no freshwater lake area. Saline lakes cover 0.02% of the State (Table 2) but they are present in only five inland catchments: Far north-west, Paroo, Warrego, Murray–Riverina and Lower Murray (Table 3). The Paroo River and Far north-west catchments cover 93% of all saline lake area (Table 3).

Coastal wetlands

Approximately 1.3% of the coastal land area is wetland (Table 2), principally estuarine wetlands (59%) and coastal lagoons and lakes (35%) (Table 2). Floodplain wetlands rank lowest in wetland area in most coastal catchments, accounting for only 5% of wetland area (Table 4). Most wetlands are less than 250 m above sea level (Fig. 3a) but are more wide spread across different climatic zones than inland wetlands (Fig. 3b).

The proportion of wetland is highest in the Macquarie–Tuggerah Lakes catchment (Table 4), and > 5% proportion in the Karuah, Georges–Cooks and Illawarra–Port Hacking catchments (Table 4). Macquarie–Tuggerah Lakes and Karuah have more than 20 000 ha of wetland area, and Clarence, Hawkesbury–Nepean, Georges–Cooks and Clyde have more than 10 000 ha of wetland area. In 14 of the 22 coastal catchments, estuarine wetlands account for 50% or more of the wetlands in the catchment and more than 80% in six of these catchments (Table 4). Coastal lagoons and lakes are concentrated in the Clyde (22) and Bega–Dry (24), with Clarence, Bellinger, Hastings, Karuah, Macquarie–Tuggerah and Deua, and Tuross with 10 or more coastal lagoons and lakes (Table 4). The Karuah and Macquarie–Tuggerah catchments have most of the area of coastal lagoons and lakes (Table 4). Floodplain wetlands are proportionally greater than 10% in the Richmond, Clarence, Macleay, Manning, Hunter and Shoalhaven (Table 4). Compared to inland, there are few freshwater lakes in the coastal catchments and they cover a small part of the land area (Table 4).

Wetland conservation

Less than 1% of the State's wetland area is listed as internationally significant under the Ramsar Convention, whereas 3% is part of National Parks and Wildlife Service (NPWS) reserves and 21% of the State's wetland area is of national importance (DIWA) (Table 2; Fig. 4). There is an additional 3% of the State's wetlands in State forests (mostly inland) and 0.7% of all wetland area that is protected under SEPP 14 (Table 2; Fig. 4). Although the largest areas of wetlands in NPWS reserves are in inland catchments, the greatest proportion of wetlands is reserved in coastal catchments (Table 2; Fig. 4).

Table 1. Spatially derived wetland groups (and reservoirs) for inland (Murray–Darling Basin and Far north-west) and coastal wetlands across New South Wales

| Global groups ^a | Categories | Description |
|----------------------------|---------------------------|--|
| Palustrine and riverine | Floodplain wetland | River and creek channels and adjacent inundated vegetation, which includes swamps, waterholes and shallow depressions. These areas are defined by the satellite imagery, which was chosen to reflect large floods over a 10-year period. |
| Estuarine | Estuarine wetland | Open water bodies and adjacent vegetation at the mouth of a river open to the sea where salt water and freshwater mix (e.g. Port Stephens). Estuarine extent was determined by tidal influences. |
| Lacustrine | Freshwater lake | Naturally occurring drainage basins of open water and not estuarine or coastal lagoons and lakes or saline on a 1:250 000 map and where surface aquatic vegetation did not dominate spectral reflectance. |
| | Saline lake | Inland naturally occurring drainage basins of saline open water, annotated as 'salt' or 'salt lake' on 1:250 000 topographic maps. |
| | Coastal lagoons and lakes | Open bodies of water and adjacent vegetation that were not obviously part of the river and were completely (e.g. Myall Lakes) or partly (e.g. Wamberal Lagoon) separated from the sea. |
| Reservoirs | | Open bodies of water usually created by a wall or levee, including reservoirs, farm dams, off-river storages, mining and quarry dams, sewage ponds, evaporation basins, canals and open basins. |

^aGlobal classification groups (Cowardin *et al.* 1977; Wilen and Bates 1995).

Table 2. Total area of wetlands and reservoirs (ha), and wetland areas in National Parks and Wildlife Service (NPWS) reserves, State forests, State Environmental Planning Policy No. 14 (SEPP 14), A Directory of Important Wetlands in Australia (DIWA), and listed under the Ramsar Convention within three regions; that is, New South Wales (NSW) (which includes the Australian Capital Territory), the inland (Murray–Darling Basin and Far north-west) and the coast (coastal catchments)

| Region ^a | Group | Area | % ^b | Number ^c | Conservation land measure | | | | | | | | Ramsar | |
|------------------------------------|---------------------------|-----------|----------------|---------------------|---------------------------|----------------|---------------|----------------|---------|----------------|---------|----------------|--------|----------------|
| | | | | | NPWS | | State forests | | SEPP 14 | | DIWA | | | |
| | | | | | Area | % ^d | Area | % ^d | Area | % ^d | Area | % ^d | Area | % ^d |
| Inland (66 580 816 ha) | Saline lake | 18 518 | <0.1 | 218 | 0 | 0 | 0 | 0 | — | — | 7816 | 0.2 | 0 | 0 |
| | Freshwater lake | 295 805 | 0.4 | 2204 | 18 726 | 0.4 | 915 | <0.1 | — | — | 160 951 | 3.7 | 1362 | <0.1 |
| | Floodplain wetland | 4 003 979 | 6.0 | — | 84 909 | 2.0 | 136 201 | 3.2 | — | — | 672 910 | 15.6 | 18 490 | 0.4 |
| | All wetlands | 4 318 302 | 6.5 | — | 103 635 | 2.4 | 137 116 | 3.2 | — | — | 841 677 | 19.5 | 19 852 | 0.5 |
| | Reservoir | 84 416 | 0.1 | 1898 | 2881 | — | 61 | — | — | — | 393 | — | 0 | 0 |
| Coastal (13 977 498 ha) | Freshwater lake | 1927 | <0.1 | 90 | 196 | 0.1 | 16 | 0.8 | 144 | 0.1 | 1429 | 0.8 | 100 | 0.1 |
| | Estuarine | 110 795 | 0.8 | — | 13 001 | 6.9 | 224 | 0.2 | 23 761 | 12.6 | 46 107 | 24.5 | 4360 | 2.3 |
| | Coastal lagoons and lakes | 65 871 | 0.5 | 194 | 17 755 | 9.4 | 44 | 0.1 | 6741 | 3.6 | 42 825 | 22.8 | 10 363 | 5.5 |
| | Floodplain wetland | 9304 | 0.1 | — | 1912 | 1.0 | 78 | 0.8 | 1386 | 0.7 | 1950 | 1.0 | 0 | 0 |
| | All wetlands | 187 897 | 1.3 | — | 32 864 | 17.5 | 362 | 0.2 | 32 032 | 17.0 | 92 311 | 49.1 | 14 823 | 7.9 |
| | Reservoir | 37 058 | 0.3 | 1158 | 512 | — | 30 | — | 45 | — | 164 | — | 140 | — |
| New South Wales (80 558 314 ha) | Saline lake | 18 518 | <0.1 | 218 | 0 | 0 | 0 | 0 | — | — | 7816 | 0.2 | 0 | 0 |
| | Freshwater lake | 297 732 | 0.4 | 2294 | 18 922 | 0.4 | 931 | <0.1 | 144 | <0.1 | 162 380 | 3.6 | 1462 | <0.1 |
| | Estuarine | 110 795 | 0.1 | — | 13 001 | 0.3 | 224 | <0.1 | 23 761 | 0.5 | 46 107 | 1.0 | 4360 | 0.1 |
| | Coastal lagoons and lakes | 65 871 | 0.1 | 194 | 17 755 | 0.4 | 44 | <0.1 | 6741 | 0.1 | 42 825 | 1.0 | 10 363 | 0.2 |
| | Floodplain wetland | 4 013 283 | 5.0 | — | 86 821 | 1.9 | 136 279 | 3.0 | 1386 | <0.1 | 674 860 | 15.0 | 18 490 | 0.4 |
| | All wetlands | 4 506 199 | 5.6 | — | 136 499 | 3.0 | 137 478 | 3.1 | 32 032 | 0.7 | 933 988 | 20.7 | 34 675 | 0.8 |
| | Reservoir | 121 474 | 0.2 | 3056 | 3393 | — | 91 | — | 45 | — | 557 | — | 140 | — |

^aLand areas are given in parentheses.^bRelative to land area in each region.^cNumbers are counts of lakes, lagoons or reservoirs and may include more than one polygon.^dRelative to wetland area within each region.

Number of freshwater lakes, saline lakes, coastal lakes and lagoons and reservoirs are given. Percentages are wetland area relative to total land area within a region or, for conservation land measures (NPWS, State forests, SEPP 14, DIWA, Ramsar), relative to wetland area within a region. Categories of conservation land measures are inclusive.




Fig. 2. Distribution of wetlands and reservoirs (grey shading) across New South Wales based on Landsat MSS satellite data (80 m pixel) from the period 1984–1993 for the inland and Landsat TM satellite data (30 m pixel) from the period September 1994–January 1995.

Despite 96% of wetland area occurring in inland catchments, only 0.5% of this area is listed under the Ramsar Convention, of which 2.4% occur in NPWS reserves, 3.2% in State forests and approximately 20% is defined as nationally important (DIWA) (Table 2; Fig. 4). In contrast, wetlands on the coast represent only 4% of the State's wetland area but nearly 8% are listed under the Ramsar Convention, of which 17.5% occur in NPWS reserves, almost 50% in the DIWA and 17% is protected under SEPP 14 (Table 2; Fig. 4). There were Ramsar-listed wetland areas in four inland catchments (i.e. Condamine–Culgoa, Gwydir, Macquarie–Bogan and Far north-west) (Table 5) and five coastal catchments (i.e. Clarence, Karuah, Hunter, Georges–Cook and Snowy) (Table 6). The highest proportions of inland wetland areas in NPWS reserves are in the Darling (5%), Paroo (4%), Macquarie–Bogan (4%) and Condamine–Culgoa (3%) (Table 5). Another three inland catchments, Warrego, Lachlan, and Far north-west, have more than 3000 ha reserved, whereas the Namoi and Lake George have none (Table 5). In contrast, all coastal catchments have some wetland area in NPWS reserves (Table 6). The Hunter catchment (57%) has the highest

proportion but the Brunswick, Richmond, Hastings, Manning, Karuah, Towamba and Snowy all have more than 20% of their wetlands reserved and Tweed, Bellinger, Macleay, Shoalhaven and Tuross have 10–20% of their wetlands reserved (Table 6). Tweed, Brunswick, Richmond, Bellinger, Macleay, Hastings, Manning, Karuah, Hunter, Shoalhaven, Deua, Bega–Dry and Towamba also have 20–47% of their wetland area protected under SEPP 14 (Table 6). Most wetland area in State forests (14–31%) is in the Murray–Riverina, Benanee and Lower Murray catchments (Table 5).

There are three inland catchments, Macquarie, Lake George and Murray–Riverina, with more than 50% of the wetland area listed as nationally important (DIWA) and a further two, Paroo and Gwydir, with 20–50% of the wetland area listed (Table 5). Most other catchments have relatively small areas listed as wetlands of national importance. The larger areas (> 30000 ha) of inland wetlands listed in the DIWA are in the Macquarie, Paroo, Murray–Riverina, Darling, Lachlan, Condamine–Culgoa and Far north-west catchments. Large areas on the coast (> 5000 ha) listed in the DIWA occur in the Clarence, Karuah, Macquarie–Tuggerah

Table 3. Areas (ha) of catchments, total wetlands of wetland groups and their proportions and reservoirs in inland New South Wales

| Catchment name ^a | Catchment Area | Total wetland Area | % ^b | Saline lake Area | % | No. ^c | Freshwater lake Area | % | No. ^c | Floodplain Area | % | Reservoir Area | No. ^c |
|------------------------------------|----------------|-----------------------|----------------|---------------------|-----|------------------|-------------------------|-----|-------------------|--------------------|------|-------------------|------------------|
| Paroo (Total) (1) | 7412 774 | 996455 | 13 | 14085 | 1 | 155 | 36754 | 4 | 23 | 945616 | 95 | 688 | 30 |
| NSW | 4115249 | 666913 | 16 | 8854 | 1 | 118 | 26300 | 4 | 18 | 631759 | 95 | 177 | 7 |
| Qld | 3297525 | 329542 | 10 | 5231 | 2 | 37 | 10454 | 3 | 5 | 313857 | 95 | 511 | 23 |
| Warrego (Total) (2) | 6354301 | 851077 | 13 | 520 | <1 | 15 | 7275 | 1 | 4 | 843282 | 99 | 1643 | 61 |
| NSW | 1138944 | 311952 | 27 | 520 | <1 | 15 | 411 | <1 | 3 | 311021 | 100 | 1000 | 9 |
| Qld | 5215357 | 539125 | 10 | 0 | 0 | 0 | 6864 | 1 | 1 | 532261 | 99 | 643 | 52 |
| Condamine-Culgoa (Total) (3) | 16359717 | 1453923 | 9 | 1071 | <1 | 1 | 16888 | 1 | 29 | 1435964 | 99 | 11037 | 340 |
| NSW | 2611171 | 738867 | 28 | 0 | 0 | 0 | 15898 | 2 | 8 | 722969 | 98 | 102 | 24 |
| Qld | 13748546 | 715056 | 5 | 1071 | <1 | 1 | 990 | <1 | 21 | 712995 | 100 | 10935 | 316 |
| Darling (4) | 11258100 | 581135 | 5 | 0 | 0 | 0 | 130730 | 22 | 140 | 450405 | 78 | 2983 | 121 |
| Border and Moonie (Total) (5) | 6299664 | 108832 | 2 | 0 | 0 | 0 | 180 | <1 | 14 | 108652 | 100 | 8457 | 191 |
| NSW | 2463285 | 60949 | 2 | 0 | 0 | 0 | 180 | <1 | 14 | 60769 | 100 | 3309 | 88 |
| Qld | 3795253 | 47883 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 47883 | 100 | 5148 | 103 |
| Gwydir (6) | 2659603 | 58907 | 2 | 0 | 0 | 0 | 115 | <1 | 2 | 58792 | 100 | 15650 | 193 |
| Namoi (7) | 4195075 | 52677 | 1 | 0 | 0 | 0 | 6902 | 13 | 4 | 45775 | 87 | 8721 | 162 |
| Castlereagh (8) | 1739480 | 16949 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 16949 | 100 | 1041 | 121 |
| Macquarie-Bogan (9) | 7463395 | 421516 | 6 | 0 | 0 | 0 | 2173 | 1 | 36 | 419343 | 99 | 13391 | 239 |
| Lachlan (10) | 9064850 | 471011 | 5 | 0 | 0 | 0 | 25217 | 5 | 24 | 445794 | 95 | 7769 | 163 |
| Murrumbidgee (11) | 8152712 | 277369 | 3 | 0 | 0 | 0 | 14436 | 5 | 33 | 262933 | 95 | 18526 | 337 |
| Lake George (12) | 94069 | 13018 | 14 | 0 | 0 | 0 | 12923 | 99 | 3 | 95 | 1 | 57 | 3 |
| Murray catchments ^d | 5081152 | 372717 | 7 | 800 | <1 | 34 | 21697 | 6 | 55 | 350220 | 94 | 11179 | 271 |
| Upper Murray (Total) (13) | 1535302 | 940 | <1 | 0 | 0 | 0 | 3 | <1 | 2 | 937 | 100 | 23535 | 54 |
| NSW | 520544 | 292 | <1 | 0 | 0 | 0 | 1 | <1 | 1 | 291 | 100 | 6967 | 26 |
| Vic | 1014758 | 648 | <1 | 0 | 0 | 0 | 2 | <1 | 1 | 646 | 100 | 16568 | 28 |
| Murray-Riverina (14) | 1502859 | 262430 | 17 | 20 | <1 | 1 | 2360 | 1 | 8 | 260050 | 99 | 3468 | 85 |
| Benanee (15) | 2129292 | 63834 | 3 | 0 | 0 | 0 | 3615 | 6 | 13 | 60219 | 94 | 0 | 0 |
| Lower Murray (Total) (16) | 5813578 | 280882 | 5 | 2449 | 1 | 42 | 127448 | 45 | 62 | 150985 | 54 | 2269 | 259 |
| NSW | 928457 | 46161 | 5 | 780 | 2 | 33 | 15721 | 34 | 33 | 29660 | 64 | 744 | 160 |
| SA | 4885121 | 234721 | 5 | 1669 | 1 | 9 | 111727 | 48 | 29 | 121325 | 52 | 1525 | 99 |
| Far north-west ^e (17) | 6543731 | 274322 | 4 | 8344 | 3 | 51 ^f | 38823 | 14 | 1864 ^g | 227155 | 83 | 511 | 160 |
| Total of all wetlands ^h | 66580816 | 4318302 | 6.5 | 18518 | 0.4 | 218 | 295805 | 6.9 | 2204 | 4003979 | 92.7 | 84416 | 1898 |

^aFor Murray-Darling Basin catchments that extend across State borders, a catchment total including separate state portions (NSW, New South Wales; Qld, Queensland; Vic, Victoria; SA, South Australia) is given. For all other catchments, areas are the NSW portions only. See Fig. 1 for locations of catchments signified by numbers in parentheses.

^bPercentages of wetland area relative to total land area within the catchment.

^cNumbers of wetlands are counts of lakes, lagoons or reservoirs and may include more than one polygon.

^dAll catchments in the Murray totalled.

^eIncludes Lake Bancannia and NSW portions of Lake Frome, Cooper Creek and Bulloo River (see Fig. 1).

^fFifty per cent of these wetlands are < 5 ha (53 ha or 0.6% of wetland area).

^gSixty-two per cent of these wetlands are < 5 ha (2836 ha or 1% of wetland area).

^hDoes not include parts of catchments within other States.

Percentage of total wetland area in each of the 17 catchments (1-17; see Fig. 1) and numbers of discrete freshwater lakes, saline lakes and reservoirs are given. Percentages for total wetland are relative to catchment land area and for wetland groups, relative to total wetland area.




Fig. 3. Percentages of wetland area across New South Wales (filled), inland (hatched) and coast (open) in relation to: (a) elevation (metres above sea level) and (b) annual rainfall. Numbers indicate the percentage of land area in each of the elevation and rainfall classes.

and Clyde catchments, and the Macleay, Karuah, Hunter, Illawarra–Hacking and Shoalhaven catchments have approximately 50% or more of their wetland area listed in the DIWA (Table 6).

Water resource development and intensive agriculture

Nine of the 17 inland catchments have more than 10% of their catchment area used for intensive agriculture (Table 5), whereas four catchments (Far north-west, Paroo, Warrego and Darling), in the north-west of the State have less than 1% of their land area utilised by intensive agriculture. Of the inland catchments, the Murray–Riverina, Gwydir, Macquarie–Bogan, Murrumbidgee, Lachlan and Border catchments have more than 20% of their catchment area used for intensive agriculture. On the coast, Macquarie–Tuggerah Lakes, Brunswick, Illawarra–Port Hacking and Tweed catchments have more than 20% of their catchment area used for intensive agriculture (Table 6).

The potential capacity of dams across the State is 29347000 ML and inland catchments account for 64% of this capacity (Tables 5,6). Nine catchments have storage capacities greater than 1000000 ML; that is, Upper Murray, Murrumbidgee, Darling, Macquarie, Lachlan, Gwydir, Snowy, Hawkesbury–Nepean and Hunter (Tables 5,6). The Macquarie, Lachlan, Murrumbidgee and Hawkesbury–Nepean catchments have more than 300 weirs, and the Murray–Riverina, Namoi, Richmond, Hunter and Clarence

have more than 100 (Tables 5,6). In the inland, the Castlereagh, Condamine–Culgoa, Far north-west, Paroo and Warrego catchments have few weirs and a low storage capacity of Government-built dams (Table 5), whereas similar catchments on the coast include the Brunswick, Bellinger, Deua, Genoa, Hastings, Illawarra–Port Hacking, Karuah, Manning, Towamba and Tuross catchments (Table 6).

Discussion

Wetland inventory and classification

Lack of data for the large-scale distribution and extent of wetlands has hampered effective wetland conservation around the world usually because governments need to set conservation priorities at large jurisdictional or catchment scales. Despite the imperative for these data, the state of wetland inventory around the world remains parlous (Finlayson *et al.* 1999). Technology in the form of remote sensing makes large-scale inventory possible, given sufficient resources, but there are two other constraints to progress in inventory development: multiple objectives and classification.

Finlayson *et al.* (1999) advocated objectives that included eight essential core data elements and six additional information categories for inventories. We achieved only two of the former (area and boundary, and location) and two of the latter (land use and conservation status) in our inventory. Of these, the latter were based on overlaying other data layers but were not collected in the present study (Tables 5,6). Collection of additional essential data (water regimen, chemistry and biota) is confounded by temporal scale, with characterisation demanding considerable resources that are seldom available for large areas. Furthermore, a variable such as ‘biota’ includes a huge range of plants, animals and communities. Although other objectives are important, they are not easily achievable on a large scale.

The second constraint of *a priori* classification can also hamper progress. Classification of wetlands adds differentiation to an audit of location and extent (Semeniuk and Semeniuk 1995, 1997) and helps analysis of representativeness or loss or degradation (Wilén and Bates 1995). Yet, it is also specific to objectives and simplifies spatial and temporal variation. A geomorphic classification differs from an ecological one and, within ecological classifications, one based on vegetation will differ from one based on an animal species. Also, limits to methodology compound problems with definition. Even for our broad classification we could not adequately and objectively differentiate between palustrine and riverine areas from satellite imagery (Table 1; Wilén and Bates 1995). This is partly because of inherent problems with the definition for the palustrine category (see Semeniuk and Semeniuk 1997). Our broad classification was *a posteriori* and based primarily on location and broad

Table 4. Areas (ha) of catchments, total wetlands, wetland groups and their proportions and reservoirs in coastal New South Wales

| Catchment name | Catchment Area | Total wetland Area | % ^a | Freshwater lake Area | % | No. ^b | Estuarine Area | % | Coastal lakes and lagoons Area | % | No. ^b | Floodplain Area | % | Reservoir Area | No. ^b |
|-------------------------------|----------------|--------------------|----------------|----------------------|----|------------------|----------------|----|--------------------------------|----|------------------|-----------------|----|----------------|------------------|
| Tweed (18) | 132561 | 3708 | 3 | 1 | <1 | 1 | 3439 | 93 | 199 | 5 | 3 | 69 | 2 | 263 | 43 |
| Brunswick (19) | 27367 | 724 | 3 | 0 | 0 | 0 | 713 | 98 | 9 | 1 | 3 | 2 | <1 | 14 | 2 |
| Richmond (20) | 703082 | 9072 | 1 | 45 | <1 | 15 | 6965 | 77 | 657 | 7 | 9 | 1405 | 15 | 423 | 60 |
| Clarence (21) | 2265802 | 16105 | 1 | 197 | 1 | 4 | 10,810 | 67 | 3173 | 20 | 15 | 1925 | 12 | 252 | 36 |
| Bellinger (22) | 305264 | 4592 | 2 | 0 | 0 | 0 | 3988 | 87 | 409 | 9 | 13 | 195 | 4 | 78 | 46 |
| Macleay (23) | 1140210 | 4965 | 0 | 34 | 1 | 3 | 2882 | 58 | 642 | 13 | 8 | 1407 | 28 | 337 | 61 |
| Hastings (24) | 448394 | 8383 | 2 | 54 | 1 | 2 | 4553 | 54 | 3693 | 44 | 12 | 83 | 1 | 92 | 20 |
| Manning (25) | 821838 | 5708 | 1 | 0 | 0 | 0 | 3822 | 67 | 1201 | 21 | 7 | 685 | 12 | 155 | 68 |
| Karuah (26) | 449803 | 44384 | 10 | 5 | <1 | 3 | 25,822 | 58 | 18424 | 42 | 11 | 133 | <1 | 147 | 62 |
| Hunter (27) | 2142494 | 5979 | 0 | 64 | 1 | 10 | 4851 | 81 | 2 | <1 | 1 | 1062 | 18 | 8632 | 270 |
| Macquarie–Tuggerah Lakes (28) | 183225 | 22768 | 12 | 20 | <1 | 3 | 3350 | 15 | 19392 | 85 | 15 | 6 | <1 | 203 | 41 |
| Hawkesbury–Nepean (29) | 2181872 | 15841 | 1 | 1086 | 7 | 10 | 13674 | 86 | 0 | 0 | 0 | 1081 | 7 | 10084 | 303 |
| Georges–Cook (30) | 182177 | 12514 | 7 | 1 | <1 | 1 | 12158 | 97 | 336 | 3 | 5 | 19 | <1 | 930 | 43 |
| Illawarra–Port Hacking (31) | 80029 | 5181 | 6 | 0 | 0 | 0 | 1686 | 33 | 3492 | 67 | 8 | 3 | <1 | 39 | 13 |
| Shoalhaven (32) | 724089 | 4465 | 1 | 1 | <1 | 1 | 3001 | 67 | 690 | 15 | 3 | 773 | 17 | 1247 | 30 |
| Clyde (33) | 328449 | 13060 | 4 | 0 | 0 | 0 | 6414 | 49 | 6633 | 51 | 22 | 13 | <1 | 98 | 9 |
| Deua (34) | 149543 | 804 | 1 | 0 | 0 | 0 | 513 | 64 | 291 | 36 | 10 | 0 | 0 | 5 | 4 |
| Tuross (35) | 215890 | 3717 | 2 | 4 | <1 | 3 | 665 | 18 | 3031 | 82 | 16 | 17 | <1 | 3 | 4 |
| Bega–Dry (36) | 284302 | 2977 | 1 | 3 | <1 | 3 | 687 | 23 | 2271 | 76 | 24 | 16 | 1 | 174 | 25 |
| Towamba (37) | 216600 | 2144 | 1 | 9 | <1 | 3 | 802 | 37 | 1326 | 62 | 9 | 7 | <1 | 27 | 6 |
| Genoa (38) | 111352 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Snowy (39) | 883155 | 806 | 0 | 403 | 50 | 28 | 0 | 0 | 0 | 0 | 0 | 403 | 50 | 13855 | 12 |
| Total of all wetlands | 13977498 | 187897 | 1.3 | 1927 | 1 | 90 | 110795 | 59 | 65871 | 35 | 194 | 9304 | 5 | 37058 | 1158 |

^aPercentages of wetland area relative to total land area within the catchment.^bNumbers of wetlands are counts of lakes, lagoons or reservoirs and may include more than one polygon.

Percentage of total wetland area in each of the 22 coastal catchments (18–39; see Fig. 1) and numbers of discrete freshwater lakes, coastal lagoons, lakes and reservoirs are given. Percentages for total wetland are relative to catchment land area, and for wetland groups are relative to total wetland area.




Fig. 4. Inland and coastal distribution of: (a) all wetlands and areas with conservation status in New South Wales; (b) National Parks and Wildlife Service (NPWS) reserves; (c) State forests; (d) State Environmental Planning Policy No. 14 (SEPP 14); (e) Directory of Important Wetlands in Australia (DIWA); and (f) wetlands listed as internationally important under the Ramsar Convention. Areas are given in parentheses. Dotted line (north–south) separates inland from coastal catchments.

geomorphic categories (Cowardin *et al.* 1979; Wilen and Bates 1995; Semeniuk and Semeniuk 1995, 1997), which were possibly less prone to problems of definition and coped with lack of data over large scales and effects of temporal changes. We showed that it was possible to develop a large-scale inventory (Fig. 2), determine the distribution and extent of those wetlands that allowed analyses of conservation priorities.

Wetland distribution and extent

Australia has no reasonable estimates of wetland area (Finlayson *et al.* 1999) and our estimate for NSW could form part of a future continental estimate. Approximately 6% coverage of land area in NSW is covered by wetlands (Table 2), which is similar to that covered in North America (Wilen and Bates 1995) but significantly lower than that covered in more mesic parts of the world such as northern

Canada (20%) and Eastern Europe (11.8%) (Stevenson and Frazier 1999). Most wetland area in NSW (83%) lies in arid regions (< 500 mm annual rainfall) (Fig. 3b) and not in humid regions; areas not well defined by national attempts to estimate wetland areas (Pajmans *et al.* 1985; Aselmann and Crutzen 1989). Freshwater lakes occur in many catchments but salt lakes are distributed primarily in the most arid regions of the State; that is, the Paroo and Far north-west catchments (Table 3).

Although occupying approximately 17% of the land area, the coastal region of NSW only has about 4% of the wetland area (Fig. 2), with relatively little floodplain wetland (Tables 2,3). Coastal catchments are considerably smaller than inland catchments (Fig. 1) and most (~60%) coastal catchments are above 250 m above sea level, the elevation below which most wetlands are located (Fig. 3a). Also, most (96%) of the mean annual run-off from rivers on the south-east coast flows out to sea, compared with only 24% of the flow in the Murray–Darling Basin in inland NSW (National Land and Water Resources Audit 2001b). Finally, drainage, alteration of the floodplain (Pressey 1989) and urbanisation (Adam 1995) have destroyed or degraded wetlands in coastal NSW. Ten of the 22 coastal catchments have more than 10% of their area under intensive agriculture (Table 6). Major population centres such as Sydney in the Hawkesbury–Nepean and Georges–Cooks catchments, Newcastle in the Hunter and Wollongong in the Shoalhaven also have large water resource developments (Table 6) that reduce flooding.

Wetland conservation

The main strategy for conservation of wetlands around the world is based on sites (Amezaga *et al.* 2002), such as National Parks and Wildlife Service (NPWS) reserves. This approach is also used to list wetlands of international importance under the Ramsar Convention, SEPP 14 or wetlands of national importance (Tables 2,5,6; Fig. 4). The overlap between NPWS reserves and wetland distribution is approximately 3% of the 4.5 million ha of wetland (Table 2; Fig. 4) and occurs disproportionately on the coast compared to inland (Tables 2,5,6; Fig. 4). Coastal catchments have up to 60% (Karuah) of their wetlands reserved (Table 6). This illustrates the problem for wetland conservation, which is focused generally on coastal systems around the world (La Peyre *et al.* 2000) where there is greater knowledge and more conservation effort. Until distribution of all wetland areas is known, conservation effort will remain biased.

Traditional conservation measures (e.g. reserves, Ramsar sites) seldom protect dependent biota and ecological processes adequately from threats operating at catchment scales (Barendregt *et al.* 1995). This is further complicated because many of the world's rivers flow between jurisdictional boundaries and so effective conservation requires integration of river management planning and

management of threats. Wetlands listed under the Ramsar Convention in Australia or the wetlands of national importance can trigger assessment of threatening processes under provisions of national conservation legislation. The building of dams, floodplain development and diversion of water resources are probably the most deleterious and pervasive threats to inland river systems (Lemly *et al.* 2000). About 89% of all surface water used in NSW (9000000 ML in 1996/1997) was for irrigation, increasing by 52% between 1983/1984 and 1996/1997 (National Land and Water Resources Audit 2001b). Our wetland inventory identifies the focus for conservation and management against potential threats (Tables 5,6). Most major regulated rivers (Murray, Murrumbidgee, Lachlan, Macquarie, Namoi, Gwydir and Border) have significant potential storage capacity and approximately 20% or more of their catchments is used for intensive agriculture, resulting in many wetland areas becoming reduced (Kingsford 2000a,b). The concentration of wetlands (46%) in four north-west catchments (Paroo, Warrego, Far north-west, Condamine–Culgoa) with relatively little water resource development should be a focus for wetland conservation and threat management. One of these, the Condamine–Culgoa, underwent considerable water resource development during the 1990s (Kingsford 2000b; Thoms and Parsons 2003), potentially affecting as much as half of the floodplain wetland identified in the NSW portion of the Condamine–Culgoa catchment (i.e. ~350000 ha). Apart from on the coast (Table 6), urban development is not a direct threat to wetlands because it is sparse (Tables 5 and 6) but, indirectly, people living on the coast have food and clothing requirements that affect the conservation of wetlands on inland rivers. The impact and threat of the draining of wetlands could not be quantified because of the lack of distributional data.

In conclusion, wetlands are among the more diverse ecosystems in the world but many continue to be destroyed or degraded by anthropogenic impacts. Mitigation of further degradation and loss is dependent partly on a strategic approach to wetland conservation and assisted by large-scale inventories. The first step is to produce a simple inventory that is not hampered by multiple objectives or disagreement about classification.

We developed such a wetland inventory for a considerable area of Australia (Fig. 2), with immediate applications. The inventory was the first step to developing priorities for conservation effort across the State and the management of threats at different scales; that is, catchments or the entire province (i.e. NSW). Much still remains unknown for wetlands of NSW, including relative conservation value; hydrology; the distribution of aquatic biota; and spatial scale of threats affecting ecological health. Our inventory can form the basis for related questions. It can also be used for other *a posteriori* classifications that relate to particular objectives (e.g. geomorphology and vegetation). Ideally, the physical,

Table 5. Areas (ha) of wetland covered by different land conservation measures (National Parks and Wildlife Service (NPWS) reserves, State forests, State Environmental Planning Policy No. 14 (SEPP 14), A Directory of Important Wetlands in Australia (DIWA) and listed under the Ramsar Convention) in each of the 17 inland catchments in New South Wales (1–17, see Fig. 1), and major threats in terms of water resource development and area covered by intensive agriculture

| Catchment | NPWS | | Land conservation measure ^a | | Ramsar | | Intensive agriculture ^c % ^d | Threat No. weirs | Storage capacity (× 10 ³ ML) ^e |
|---|--------|-----|--|-------------------|--------|------|---|---------------------|---|
| | Area | % | State forests | DIWA ^b | Area | % | | | |
| | | | Area | % | Area | % | | | |
| Paroo (1) | 26605 | 4 | 0 | 0 | 181962 | 27 | 0 | 0 | 0 |
| Warrego (2) | 3036 | 1 | 0 | 0 | 1095 | 0 | 0 | 0.8 | 42 |
| Condamine–Culgoa (3) | 18527 | 3 | 0 | 0 | 37716 | 5 | 3161 | 10.4 | 27 |
| Darling (4) | 29237 | 5 | 0 | 0 | 82755 | 14 | 0 | 0 | 1896 |
| Border (Moonie) ^f (5) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 22.2 (14.6) | 67 |
| Gwydir (6) | 82 | <1 | 0 | 0 | 28304 | 48 | 708 | 1 | 1368 |
| Namoi (7) | 0 | 0 | 0 | 0 | 6776 | 13 | 0 | 0 | 880 |
| Castlereagh (8) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 20.0 | 37 |
| Macquarie (9) | 15183 | 4 | 125 | 0 | 218326 | 52 | 15743 | 4 | 1716 |
| Lachlan (10) | 3268 | <1 | 5766 | 1 | 78584 | 17 | 0 | 0 | 323 |
| Murrumbidgee (11) | 146 | <1 | 14676 | 5 | 7481 | 3 | 0 | 0 | 4206 |
| Lake George (12) | 0 | 0 | 0 | 0 | 12833 | 99 | 0 | 0 | — |
| Murray catchments | 903 | <1 | 116549 | 31 | 150863 | 40 | 0 | 0 | 6613 |
| Upper Murray (13) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 5652 |
| Riverina (14) | 0 | 0 | 95725 | 36 | 148879 | 57 | 0 | 0 | 206 |
| Benanee (15) | 903 | 1 | 14223 | 22 | 1964 | 3 | 0 | 0 | 75 |
| Lower Murray (16) | 0 | 0 | 6601 | 14 | 20 | <1 | 0 | 0 | 680 |
| Far north-west (Lake Bannania) ^f (17) | 6648 | 2 | 0 | 0 | 34982 | 13 | 240 | <1 | 8 |
| Total of all wetlands | 103635 | 2.4 | 137116 | 3.2 | 841677 | 19.5 | 19852 | <1 | 18751 |

^aConservation measure areas are mutually inclusive.

^bA Directory of Important Wetlands in Australia (Environmental Australia# 2001).

^cNational Land and Water Resources Audit (2001b) for catchments that cross the New South Wales border. Proportions include areas outside the state border.

^dPercentage relative to catchment land area.

^eDerived from Kingsford (1995) and National Land and Water Resources Audit (2001b) and where storage capacities differ, the larger storage capacity is used. Interstate catchments storage capacities include areas outside New South Wales.

^fSeparate estimates for agricultural land proportion only are given in parentheses.

Percentages are relative to total wetland area within a catchment.

Table 6. Areas (ha) of wetland covered by different conservation land measures (National Parks and Wildlife Service (NPWS) reserves, State forests, State Environmental Planning Policy No. 14 (SEPP 14), A Directory of Important Wetlands in Australia (DIWA), and listed under the Ramsar Convention) in each of the 22 coastal catchments in New South Wales (18–39; see Fig. 1), and major threats in terms of water resource development and area covered by intensive agriculture

| Catchment | Conservation land measure ^a | | | | | | | | | | Threat No. weirs | Storage capacity (× 10 ³ ML) ^d | |
|-------------------------------|--|------|---------------|-----|----------------------|------|-------------------|------|--------|-----|---------------------|--|---|
| | NPWS | | State forests | | SEPP 14 ^b | | DIWA ^c | | Ramsar | | | | Intensive agriculture ^d % ^e |
| | Area | % | Area | % | Area | % | Area | % | Area | % | | | |
| Tweed (18) | 689 | 19 | 0 | 0 | 841 | 23 | 853 | 23 | 0 | 0 | 22.8 | 36 | 17 |
| Brunswick (19) | 171 | 24 | 0 | 0 | 294 | 41 | 0 | 0 | 0 | 0 | 30.8 | 7 | 0 |
| Richmond (20) | 2370 | 26 | 62 | 1 | 3506 | 39 | 1787 | 20 | 0 | 0 | 9.9 | 188 | 34 |
| Clarence (21) | 1399 | 9 | 0 | 0 | 1943 | 12 | 11502 | 71 | 59 | <1 | 5.1 | 106 | 7 |
| Bellinger (22) | 556 | 12 | 45 | 1 | 1513 | 33 | 0 | 0 | 0 | 0 | 11.9 | 23 | 0 |
| Macleay (23) | 904 | 18 | 34 | 1 | 1430 | 29 | 2452 | 49 | 0 | 0 | 14.3 | 42 | 19 |
| Hastings (24) | 3527 | 42 | 0 | 0 | 3333 | 40 | 2975 | 35 | 0 | 0 | 3.7 | 48 | 2 |
| Manning (25) | 1147 | 20 | 0 | 0 | 2134 | 37 | 1758 | 31 | 0 | 0 | 6.1 | 27 | 4 |
| Karuah (26) | 12973 | 29 | 170 | <1 | 8962 | 20 | 35249 | 79 | 11224 | 25 | 6.6 | 11 | 0 |
| Hunter (27) | 3388 | 57 | 0 | 0 | 2807 | 47 | 3873 | 65 | 3099 | 52 | 12.1 | 123 | 1670 |
| Macquarie–Tuggerah Lakes (28) | 168 | 1 | 0 | 0 | 691 | 3 | 9516 | 42 | 0 | 0 | 34.0 | 49 | 89 |
| Hawkesbury–Nepean (29) | 1494 | 9 | 0 | 0 | 98 | 1 | 1069 | 7 | 0 | 0 | 19.0 | 444 | 2929 |
| Georges–Cook (30) | 443 | 4 | 0 | 0 | 0 | 0 | 974 | 8 | 400 | 3 | 8.5 | 47 | 124 |
| Illawarra–Port Hacking (31) | 132 | 3 | 0 | 0 | 225 | 4 | 3702 | 71 | 0 | 0 | 23.1 | 25 | 1 |
| Shoalhaven (32) | 819 | 18 | 0 | 0 | 1129 | 25 | 3486 | 78 | 0 | 0 | 13.7 | 48 | 178 |
| Clyde (33) | 981 | 8 | 0 | 0 | 1448 | 11 | 9223 | 71 | 0 | 0 | 1.6 | 15 | 7 |
| Deua (34) | 61 | 8 | 0 | 0 | 285 | 35 | 76 | 9 | 0 | 0 | 3.5 | 6 | 5 |
| Tuross (35) | 571 | 15 | 21 | 1 | 341 | 9 | 1126 | 30 | 0 | 0 | 7.5 | 8 | 0 |
| Bega–Dry (36) | 161 | 5 | 30 | 1 | 543 | 18 | 1324 | 44 | 0 | 0 | 9.1 | 56 | 16 |
| Towamba (37) | 639 | 30 | 0 | 0 | 509 | 24 | 1023 | 48 | 0 | 0 | 1.6 | 5 | 0 |
| Genoa (38) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1.5 | 0 | 0 |
| Snowy (39) | 271 | 34 | 0 | 0 | 0 | 0 | 343 | 43 | 41 | 5 | 11.8 | 24 | 5494 |
| Total of all wetlands | 32864 | 17.5 | 362 | 0.2 | 32032 | 17.0 | 92311 | 49.1 | 14823 | 7.9 | | 1338 | 10596 |

^aConservation measure areas are mutually inclusive.

^bState Environmental Planning Policy No. 14.

^cA Directory of Important Wetlands in Australia (Environment Australia 2001).

^dDerived from National Land and Water Resources Audit (2001a, 2001b) and Kingsford (1995) and where storage capacities differ, the larger storage capacity is used.

^ePercentage relative to catchment land area.

Percentages are relative to total wetland area within a catchment.

chemical or biotic data layers overlaid on an inventory of the distribution and extent of wetlands can derive objective classifications for biota (Munger *et al.* 1998) and landscapes (Gwin *et al.* 1999). Inventory information for wetlands allows priorities to be set for conservation but this is not a panacea. Further loss or degradation of wetlands will only be avoided with political, community and bureaucratic commitment to their conservation.

Acknowledgments

Funding to map the coastal wetlands and the far west of NSW was provided through the Natural Heritage Trust Fund, and funding to map the wetlands in the Murray–Darling Basin was provided by the Murray–Darling Basin Commission and NSW National Parks and Wildlife Service. We thank P. Adam and B. Pressey for their comments, which greatly improved this manuscript. We thank the National Land and Water Resources Audit for permission to use their data.

References

- Adam, P. (1995). Urbanization and transport. In 'Conserving Biodiversity: Threats and Solutions'. (Eds R. Bradstock, T. D. Auld, D. A. Keith, R. T. Kingsford, D. Lunney and D. Sivertsen.) pp. 55–75. (Surrey Beatty and Sons: Sydney, Australia.)
- Amezaga, J. M., Santamaria, L., and Green, A. J. (2002). Biotic wetland connectivity – supporting a new approach for wetland policy. *Acta Oecologica* **23**, 213–222. doi:10.1016/S1146-609X(02)01152-9
- Aselmann, I., and Crutzen, P. J. (1989). Global distribution of natural freshwater wetlands and rice paddies, their net primary productivity, seasonality and possible methane emissions. *Journal of Atmospheric Chemistry* **8**, 307–358.
- AUSLIG (1994). 'GEODATA TOPO-250K (Series 1) Topographic Data.' (Geoscience Australia: Canberra, Australia.)
- Barendregt, A., Wassen, M. J., and Schot, P. P. (1995). Hydrological systems beyond a nature reserve, the major problem in wetland conservation of Naardermeer (The Netherlands). *Biological Conservation* **72**, 393–405. doi:10.1016/0006-3207(94)00101-U
- Congalton, R. G. (1991). A review of assessing the accuracy of classification of remotely sensed data. *Remote Sensing of Environment* **37**, 35–46. doi:10.1016/0034-4257(91)90048-B
- Congalton, R. G. (2001). Accuracy assessment and validation of remotely sensed and other spatial information. *International Journal of Wildland Fire* **10**, 321–328. doi:10.1071/WF01031
- Congalton, R. G., and Green, K. (1999). 'Assessing the Accuracy of Remotely Sensed Data: Principles and Practices'. (Lewis Publishers, CRC Press Inc.: Washington, D.C.)
- Cowardin, L. M., Carter, V., Golet, F. C., and LaRoe, E. T. (1979). 'Classification of Wetlands and Deepwater Habitats of the United States.' FWS/OBS – 79/31. (US Fish and Wildlife Service: Washington, DC, USA.)
- Environment Australia (2001). 'A Directory of Important Wetlands in Australia.' 3rd Edn. (Environment Australia: Canberra, Australia.)
- Environmental Systems Research Institute, Inc. (2000). 'ARC® Version 8.0.2.' (ESRI, Inc.: Redlands, CA, USA.)
- ERDAS (2001). 'ERDAS® Imagine 8.5.' (ERDAS® Inc.: Atlanta, GA, USA.)
- Farrier, D., Lyster, R., and Pearson, L. (1999). 'The Environmental Law Book.' 3rd Edn. (Legal Centre Publishing: Sydney, Australia.)
- Finlayson, C. M., and Rea, N. (1999). Reasons for the loss and degradation of Australian wetlands. *Wetlands Ecology and Management* **7**, 1–11. doi:10.1023/A:1008495619951
- Finlayson, C. M., Davidson, N. C., Spiers, A. G., and Stevenson, N. J. (1999). Global wetland inventory – current status and future priorities. *Marine and Freshwater Research* **50**, 717–727.
- Frazier, P., and Page, K. (2000). Water body detection and delineation with Landsat TM data. *Photogrammetric Engineering and Remote Sensing* **66**, 1461–1468.
- Gwin, S. E., Kentula, M. E., and Shaffer, P. W. (1999). Evaluating the effects of wetland regulation through hydrogeomorphic classification and landscape profiles. *Wetlands* **19**, 477–489.
- Johnston, R. M., and Barson, M. M. (1993). Remote sensing of Australian wetlands: an evaluation of Landsat TM data for inventory and classification. *Australian Journal of Marine and Freshwater Research* **44**, 235–252.
- Kingsford, R. T. (1995). Ecological effects of river management in New South Wales. In 'Conserving Biodiversity: Threats and Solutions'. (Eds R. Bradstock, T. D. Auld, D. A. Keith, R. T. Kingsford, D. Lunney and D. Sivertsen.) pp. 144–161. (Surrey Beatty and Sons: Sydney, Australia.)
- Kingsford, R. T. (2000a). Ecological impacts of dams, water diversions and river management on floodplain wetlands in Australia. *Austral Ecology* **25**, 109–127. doi:10.1046/J.1442-9993.2000.01036.X
- Kingsford, R. T. (2000b). Protecting or pumping rivers in arid regions of the world? *Hydrobiologia* **427**, 1–11. doi:10.1023/A:1004033915662
- Kingsford, R. T., Wong, P. S., Braithwaite, L. W., and Maher, M. T. (1999). Waterbird abundance in eastern Australia, 1983–1992. *Wildlife Research* **26**, 351–366.
- Kingsford, R. T., Thomas, R. F., and Curtin, A. L. (2001). Conservation of wetlands in the Paroo and Warrego catchments in arid Australia. *Pacific Conservation Biology* **7**, 21–33.
- La Peyre, M. K. G., Reams, M. A., and Mendelssohn, I. A. (2000). State wetland protection: A matter of context? *Coastal Management* **28**, 287–302. doi:10.1080/08920750050133557
- Lemly, A. D., Kingsford, R. T., and Thompson, J. R. (2000). Irrigated agriculture and wildlife conservation: conflict on a global scale. *Environmental Management* **25**, 485–512. doi:10.1007/S002679910039
- Munger, J. C., Gerber, M., Madrid, K., Carroll, M. A., Petersen, W., and Heberger, L. (1998). US National Wetland Inventory classifications as predictors of the occurrence of Columbia spotted frogs (*Rana luteiventris*) and Pacific treefrogs (*Hyla regilla*). *Conservation Biology* **12**, 320–330. doi:10.1046/J.1523-1739.1998.97012.X
- National Land and Water Resources Audit. (2001a). 'Australian Agriculture Assessment 2001.' (Commonwealth of Australia: Canberra, Australia.)
- National Land and Water Resources Audit. (2001b). 'Australian Water Resources Assessment 2000.' (Commonwealth of Australia: Canberra, Australia.)
- Paijmans, K., Galloway, R. W., Faith, D. P., Fleming, P. M., Haantjens, H. A., Heyligers, P. C., Kalma, J. D., and Löffler, E. (1985). 'Aspects of Australian Wetlands.' CSIRO Division of Water & Land Resources, Technical Paper No.44. (CSIRO: Canberra, Australia.)
- Pressey, R. L. (1989). Wetlands of the Lower Clarence Floodplain, Northern Coastal New South Wales. *Proceedings of the Linnean Society of New South Wales* **111**, 143–155.
- Pressey, R. L., and Adam, P. (1995). A review of wetland inventory and classification in Australia. *Vegetatio* **118**, 81–101.
- Rempel, R. S., Abraham, K. F., Gadawski, T. R., Gabor, S., and Ross, R. K. (1997). A simple wetland habitat classification for boreal forest waterfowl. *The Journal of Wildlife Management* **61**, 746–757.
- Richards, J. A. (1993). 'Remote Sensing Digital Image Analysis. An Introduction.' 2nd Edn. (Springer-Verlag: Berlin, Germany.)
- Rosenfield, G. H., and Fitzpatrick-Lins, K. (1986). A coefficient of agreement as a measure of thematic classification accuracy. *Photogrammetric Engineering and Remote Sensing* **52**, 223–227.

- Semeniuk, C. A., and Semeniuk, V. (1995). A geomorphic approach to global classification for inland wetlands. *Vegetatio* **118**, 103–124.
- Semeniuk, V., and Semeniuk, C. A. (1997). A geomorphic approach to global classification for natural inland wetlands and rationalization of the system used by the Ramsar Convention – a discussion. *Wetlands Ecology and Management* **5**, 145–158. doi:10.1023/A:1008207726826
- Stevenson, N., and Frazier, S. (1999). 'Review of Wetland Inventory Information in Eastern Europe.' (Wetlands International: Wageningen, The Netherlands.)
- The Ramsar Convention Bureau (1999). 'The Ramsar Convention on Wetlands: The Criteria for Identifying Wetlands of International Importance.' (The Ramsar Convention Bureau: Gland, Switzerland.) Available at: www.ramsar.org/key_criteria.htm
- Thoms, M., and Parsons, M. (2003). Identifying spatial and temporal patterns in the hydrological character of the Condamine–Balonne River, Australia, using multivariate statistics. *River Research and Applications* **19**, 443–457.
- Wilén, B. O., and Bates, M. K. (1995). The US fish and Wildlife Service's National Wetlands Inventory project. *Vegetation* **118**, 153–169.

Manuscript received 27 May 2003; revised 9 October, 4 November 2003; accepted 10 November 2003.